

**Veterinary Feed Directive for Fish  
Romet 30®  
(sulfadimethoxine and ormetoprim)**

VFD Form Sequential ID, if  
appropriate

Veterinarian: \_\_\_\_\_  
Address: \_\_\_\_\_  
Phone #: \_\_\_\_\_  
FAX or email: (optional) \_\_\_\_\_

Client: \_\_\_\_\_  
Business or Home Address: \_\_\_\_\_  
Phone #: \_\_\_\_\_  
FAX or email: (optional) \_\_\_\_\_

Drug level:

- 1% Feed Rate: Sulfadimethoxine 3780 g/ton; Ormetoprim 756 g/ton
- 2% Feed Rate: Sulfadimethoxine 1890 g/ton; Ormetoprim 378 g/ton
- 3% Feed Rate: Sulfadimethoxine 1260 g/ton; Ormetoprim 252 g/ton
- 4% Feed Rate: Sulfadimethoxine 945 g/ton; Ormetoprim 189 g/ton
- 5% Feed Rate: Sulfadimethoxine 756 g/ton; Ormetoprim 151 g/ton
- 6% Feed Rate: Sulfadimethoxine 630 g/ton; Ormetoprim 126 g/ton

To provide 50 mg of active ingredient per kg (23 mg/lb) of live body weight of fish per day. Romet 30 contains 113.5 g/lb sulfadimethoxine and 22.7 g/lb ormetoprim.

Duration of Use: Feed for 5 days

**Indications for Use:**

Salmonids: For the control of furunculosis in freshwater-reared salmonids (trout and salmon) caused by *Aeromonas salmonicida* strains susceptible to sulfadimethoxine and ormetoprim combination.

Catfish: For the control of enteric septicemia of catfish caused by *Edwardsiella ictaluri* strains susceptible to sulfadimethoxine and ormetoprim combination.

\*\*\*\*\*  
**USE OF FEED CONTAINING THIS VETERINARY FEED DIRECTIVE (VFD) DRUG IN A MANNER OTHER THAN AS DIRECTED ON THE LABELING (EXTRALABEL USE) IS NOT PERMITTED.**

Approximate number of **FISH** to be treated: \_\_\_\_\_


Premise or Location of animals: \_\_\_\_\_

Special Instructions and/or other animal identifications:

**Affirmation of Intent** (for combination VFD drugs):

This VFD only authorizes the use of the VFD drug(s) cited in this order and is not intended to authorize the use of such drug(s) in combination with any other animal drugs.

**CAUTION:** If fish show no improvement within 2 to 3 days, or if signs of disease reappear after termination of treatment, reevaluate management practices, diagnosis of outbreak, and establish susceptibility of the bacterial isolate(s) to the drug.



**WARNING:** In catfish, withdraw 3 days before slaughter or release as stocker fish. In salmonids, withdraw 42 days before slaughter or release as stocker fish.



Date of VFD Issuance: \_\_\_\_\_ (dd/mm/yyyy)

Date of VFD Expiration: \_\_\_\_\_ (dd/mm/yyyy)  
(Cannot exceed 6 months after issuance)

Veterinarian's signature: \_\_\_\_\_